


Lentivirus **10X Titer-Up**

Name:	Lentivirus 10X Titer-Up
Cat. #:	P906, P906S
Application:	Increase lentivirus titer for ~ 10 times in virus packaging procedure. This product is for research use only.
Product Size:	P906: 1 mL (for packaging 500 mL lentivirus soup); P906S: 0.1 mL
Product Description:	This product is a novel recipe of small molecules designed for effective lentivirus packaging. <ul style="list-style-type: none">✓ Increase virus titer by 10 times✓ Increase viral RNA transcription✓ Increase virus particle packaging efficiency
Shipping / Storage:	Ship at room temperature and store in 4°C
Shelf Life:	12 months
Remark:	Each lot of 10X Titer-Up reagent is functionally tested in virus production experiment using 293T cells. Follow the recommended NIH BSL-2 guidelines for all materials containing Lentivirus.



HEK293 cells transduced by GFP lentivirus which was packaged with or without **10X Titer-Up**

Protocol (example of producing virus in **100mm petri dish**)

Day 1: Plating cells

1. 24 hours before transfection, coat 100mm dishes with **6 mL 1x Gelatin** for 60 min.
2. Aspirate gelatin, and plate **~5 X 10⁶** fast growing 293T cells per plate, in 10 ml medium.

Note: Before plating, pipet the 293T cell suspension vigorously to achieve **single cells**.
Plate the cells in **evenly** distribution manor.

Day 2: Transfection (using 101Bio “293T Transfection Reagent”, Cat. #: P902.)

1. **Change medium: 2 hours before** transfection, remove culture medium and add **10 mL fresh** complete culture medium (with 10% serum / **without antibiotics**)
2. In **tube 1** add: ~20 µg **DNA** (virus vector and packaging mix [101Bio, Cat. #: P904P])
 500 µL **DMEM** (serum-free, High Glucose)

 Pipet up and down to mix well
3. In **tube 2** Add: 40 µL **293T Transfection Reagent for Lentivirus Packaging**
 500 µL **DMEM** (serum-free, High Glucose)

 Gently pipet up and down to mix well
4. Incubate at room temperature (20 ~ 25°C) for 3 min.
5. Add **tube 2** into **tube 1**, pipet up and down several times. **Vortex for 10 seconds.**
6. Incubate for **15 minutes** at **room temperature.**
7. Add the incubated mixture **drop-wise** to the cells, and **gently rock / swirl** the plate.
8. Return the cells to 37°C incubator with 5% CO₂.

Day 3: Add 10X Titer-Up

Add 20 µl of 10X Titer-Up (500x) to the medium. Return the plates to the cell culture incubator.

Day 4: Collect virus

1. Collect virus supernatant twice at **48 and 72 hours** post transfection into a 50mL sterile FALCON tube. Centrifuge at **3,000 rpm** for **15 minutes** at 4°C to remove cell debris. Filter the clear supernatant through **0.45 µm** syringe tip filter.
2. The filtered clear supernatant is virus soup. Use it immediately or aliquot into sterile 1.5-mL tubes and store at **-80°C**, for **up to 3 months.**
3. (Optional) To concentrate virus, add 1 volume of **5X Lentivirus Concentration Solution** (101Bio, Cat. #: P904C) to 4 volume of the viral supernatant (volume of Lentivirus Concentration Solution vs. volume of viral supernatant = 1:4) and mix thoroughly. Put the mixture to 4°C refrigerator overnight and spin the virus pellet down next day. Please refer our P904C user manual for details.

Note: No side effect 10X Titer-Up reagent on gene expression has been detected when directly transduce 293T cells. It may be various on different cell types (cell lines). A pilot test on the side effect of 10X Titer-Up is advised on sensitive cells.

– The end –